

## Low Permeability of Core-forming Melts in Olivine Spinel

W G Minarik, F J Ryerson (Earth Sciences, L-202, Lawrence Livermore National Laboratory, Livermore, CA 94551; e-mail: minarik@llnl.gov)

Separation of the metallic core from the silicate mantle of a planetary body by porous flow of Fe-Ni-S-O melt through a solid matrix is viewed as unlikely because metal-rich melts, due to their high melt-solid interfacial energy, do not wet silicate grains at low pressures, precluding formation of an interconnected melt at low melt fraction, and leading to isolated melt pockets with a permeability of zero. At higher pressures increased solubility of oxygen in the metal-sulfide melt may reduce its interfacial energy against silicate minerals and allow the metal-sulfide melt to interconnect.

To address this suggestion, we have performed experiments at pressures up to 9 GPa to determine the microstructure of metallic melts in an olivine matrix. All melts with less than 50% S+O remain non-interconnected and there is no visible trend towards a reduced interfacial angle with increasing pressure. An experiment was performed at 14 GPa and 1500°C to test whether Fe-Ni-S-O melts wet high-pressure olivine spinel. 10 GPa eutectic composition Fe-Ni sulfide was added to <38 micron olivine along with (Mg,Fe) oxide to ensure oxygen saturation in the melt. This mixture was encapsulated in a diamond capsule during the 5 hour multi-anvil run.

Sulfide remained distributed throughout the charge both as inclusions in the silicate spinel and at grain corners. The approx. 7 vol. % sulfide melt showed no tendency to gravitationally segregate. Neither the increase in oxygen solubility with pressure nor the modified crystal structure are sufficient to create a wetting melt that can percolate towards the core along grain boundaries.

These results imply that separation of core from mantle must occur via silicate liquid/metal sulfide liquid separation at high temperatures in a magma ocean, either in smaller bodies that then accreted as differentiated bodies or as separation within an undifferentiated Earth close to its present size.

Work performed under the auspices of the U.S. D.O.E. by LLNL under contract W-7405-Eng-48.